

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Then $(1+.04t)(74200) = [(1+.08(t+\frac{1}{2})](68900)$. $... 68900 \times .08t = 5300(1+.04t)$ t = 1 year. ... time was January 1, 1895. F = (74200)(1.04) = \$77168.

II. Solution by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

Ler x=the number of days the first note had to run, and x+180 the numthe second had to run; regarding 360 days to the year.

Then we have, since the face of each note was the same,

$$74200 \left(1 + \frac{4}{100} \frac{x}{360}\right) = 68900 \left(1 + \frac{8}{100} \frac{x + 180}{360}\right).$$

Whence, dividing by 100 and transposing,

$$53 = \frac{5512x + 992160 - 2968x}{36000}$$

or, 2544x = 53(36000) - 992160 = 915840.

x=360 days, or date of each note was January 1st, 1895.

The face value must have been \$74200+the discount, which is 4% of \$74200 or \$2968; that is \$77168.

119. Proposed by G. B. M. ZERR, A.M., Ph.D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

The cost of an article is \$1 $\frac{9}{.100}$. The selling price is \$1,000. What is the gain per cent.?

Solution by C. C. BEBOUT, Professor of Mathematics, High School, Eigin, Ill., and J. M. HOWIE, Professor of Mathematics, The Nebraska State Normal School, Peru, Neb.

I do not think that the solution of Arithmetic problem 112, published in the August-September Monthly is correct. My solution of No. 119 will show wherein I differ. C. C. Bebout.

\$1.
$$\frac{9}{.001}$$
=\$1. +\$ $\frac{9}{.001}$ =\$1.+\$9000=\$9001, the selling price.

If the article costs \$9001., and sells for \$1000, there is no gain, but a loss of \$8001, which is 88.89+% of \$9001.

The question seems to be as to the meaning of the decimal point following the 1 in the expression $1.\frac{9}{.001}$. A common fraction can not have place value, nor can it give place value as a digit in a number. It is simply added to the number to which it is attached and is a fraction of the unit to which it is attached. If it stands alone it is a fraction of the understood (or named) unit. We would not write $3\frac{1}{2}4$ for 354, nor $2\frac{1}{2}$. for 25. Also $4\frac{1}{2}=4+\frac{1}{2}=4+\frac{1}{2}$. Every in-

teger in our decimal notation is, in theory, followed by a decimal point, and it can make no difference in the meaning of the expression whether the decimal point is expressed or understood. 4.2 must equal 4.5 and not 4.05, which is the equivalent of 4.02.

So
$$1.\frac{9}{.001} = 1\frac{9}{.001} = 1 + \frac{9}{.009}$$
.

Also solved by COOPER D. SCHMITT, D. A. LEHMAN, ELMER SCHUYLER, and the PROPOSER. These contributors agree that the result is 10887%. To my mind, the solution and discussion of the problem as published above are correct. Ed. F.

ALGEBRA.

94. Proposed by J. W. YOUNG, Columbus, Ohio.

Solve:
$$\left[\frac{x^2+14x+1}{p^4+14p^2+1}\right]^3 = \frac{x(x-1)^4}{p^2(p^2-1)^4}$$
.

I. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Let
$$p^2(p^2-1)^4 = A$$
, $(p^4+14p^2+1)^3 = B$.

$$\therefore Ax^{6} + 42Ax^{5} + 591Ax^{4} + 2828Ax^{3} + 591Ax^{2} + 42Ax + A$$

$$= Bx^{5} - 4Bx^{4} + 6Bx^{3} - 4Bx^{2} + Bx.$$

$$Ax^{6} + (42A - B)x^{5} + (591A + 4B)x^{4} + (2828A - 6B)x^{3} + (591A + 4B)x^{2} + (42A - B)x + A = 0.$$

$$A[x^3 + (1/x^3)] + (42A - B)[x^2 + (1/x^2)] + (591A + 4B)[x + (1/x)] + (2828A - 6B) = 0.$$

$$A[x+(1/x)]^3+(42A-B)[x+(1/x)]^2+(588A+4B)[x+(1/x)]$$
$$+(2744A-4B)=0.$$

Let
$$[x+(1/x)]=y$$
.

$$\begin{array}{l} \cdots p^2(p^2-1)^4y^3-(p^{12}+759p^8+2576p^6+759p^4+1)y^2\\ +4(p^{12}+189p^{16}+3p^8+3710p^6+3p^4+189p^2+1)y\\ -4(p^{12}-644p^{16}+3335p^8-1288p^6+3335p^4-644p^2+1)=0. \end{array}$$

Let $a=p^2+1/p^2$.

$$(2-a)^2y^3 - (a^3 + 756a + 2576)y^2 + 4(a^3 + 189a^2 + 3332)y - 4(a^3 - 644a^2 + 3332a) = 0.$$

$$(y-a)[(2-a)^2y^2-4(a^2+188a+644)y+4(a^2-644a+3332)]=0.$$